

METHODS FOR EVALUATING THE ABILITY TO METABOLIZE PHARMACEUTICALS AND COMPOSITIONS THEREFOR

REFERENCE TO RELATED APPLICATION

5 This application is a continuation-in-part application of U.S. Patent application Serial No. 09/144,367, filed August 31, 1998, which is hereby incorporated by reference in its entirety. This application claims the benefit of Provisional Patent Application Serial No. 60/271,630 filed in the U.S. Patent and Trademark Office on February 26, 2001, which is hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION

10 The present invention relates to polymorphisms in the cytochrome P450 3A4, cytochrome P450 3A5 and GSTM1 genes and methods for genotyping and phenotyping individuals for such polymorphisms.

BACKGROUND OF THE INVENTION

15 Cytochrome P450 enzymes are a heme-containing family that play central roles in oxidative, peroxidative and reductive metabolism of numerous endogenous and exogenous compounds, including many pharmaceutical agents. Substances known to be metabolized by P450 enzymes include steroids, bile acids, fatty acids, prostaglandins, leukotrienes, biogenic amines, retinoids, lipid hydroperoxides, phytoalexins,
20 pharmaceuticals, environmental chemicals and pollutants. P450 substrates also include natural plant products involved in flavor, odor and flower color. Many P450 enzymes also have functions in maintaining steady-state levels of endogenous ligands involved in ligand-modulated transcription of genes effecting growth, apoptosis, differentiation, cellular homeostasis, and neuroendocrine functions. The metabolism of foreign
25 chemicals by P450 enzymes can produce toxic metabolites, some of which have been implicated as agents responsible for birth defects and tumor initiation and progression.

The P450 gene superfamily is likely to have evolved from an ancestral gene present before the prokaryote/eukaryote divergence. The number of individual P450 genes in any mammalian species is estimated at 60 to 200. The cytochrome P450 (CYP) 3A subfamily is unique in that it is present in large amounts in human liver microsomes, and there are many forms in the subfamily. Several human cDNAs encoding CYP3A proteins have been identified. The most important of these are CYP3A4, CYP3A5 and CYP3A7. CYP3A4 and CYP3A7 genes are 87% homologous by amino acid and 95% homologous by nucleotide sequence, while CYP3A4 and CYP3A5 are only 88% homologous in the coding region. CYP3A4 and CYP3A7 are 91% homologous in the 5'-flanking sequences, differing by the presence of a unique P450NF specific element (NFSE) and a P450HFLa specific element (HFLaSE), respectively (Hashimoto et al, 1993).

Genetic polymorphisms of cytochrome P450 enzymes result in subpopulations of individuals that are distinct in their ability to perform particular drug biotransformation reactions. These phenotypic distinctions have important implications for selection of drugs. For example, a drug that is safe when administered to the majority of humans may cause intolerable side-effects in an individual suffering from a defect in a cytochrome P450 enzyme required for detoxification of the drug. Alternatively, a drug that is effective in most humans may be ineffective in a particular subpopulation because of the lack of a particular cytochrome P450 enzyme required for conversion of the drug to a metabolically active form. Accordingly, it is important for both drug development and clinical use to screen drugs to determine which cytochrome P450 enzymes are required for activation and/or detoxification of the drug.

It is also important to identify those individuals who are deficient in a particular P450 enzyme. This type of information has been used to advantage in the past for developing genetic assays that predict phenotype and thus predict an individual's ability to metabolize a given drug. Information such as this would be of particular value in determining the likely side effects and therapeutic failures of various drugs and routine phenotyping could be recommended for certain categories of patients.

The CYP3A subclass catalyzes a remarkable number of oxidation reactions of clinically important drugs such as quinidine, warfarin, erythromycin, cyclosporin A, midazolam, lidocain, nifedipine, and dapson. Current estimates are that more than 60% of clinically used drugs are metabolized by the CYP3A4 enzyme, including such major drug classes as calcium channel blockers, immunosuppressors, macrolide antibiotics and anticancer drugs, see Brian *et al.*, 1990, Biochemistry, vol. 29, pages 11280-11292.

Expression profiles for each member of this family varies significantly. CYP3A4 is expressed in all adult human liver and intestine, accounting for more than 50% of total P450 in both organs. Expression is inducible *in vivo* and *in vitro* by numerous compounds such as rifampicin, barbiturates and glucocorticoids. In kidney, CYP3A4 is expressed polymorphically. CYP3A4 expression is sex-influenced, as females have 24% greater expression than males. Substantial inter-individual variation in the metabolism of specific compounds by CYP3A4 has been reported (Kleinbloesem *et al.*, Biochemical Pharmacology, 1984, vol. 33, pages 3721-3724. U.S. Patent No. 6,174,684 to Rebbeck *et al.* discloses a CYP3A4 variant associated with a heightened risk of developing or having prostate cancer and a decreased risk for developing treatment-related leukemias. The polymorphism disclosed by Rebbeck *et al.* is an A to G transition in the promoter region of the CYP 3A4 gene which is thought to alter the nifedipine-specific binding element located 287 to 296 bases 5' to the CYP 3A4 transcription start site. The genotype associated with this variant is believed to increase the production of potentially DNA damaging reactive intermediates upon patient exposure to an epipodophyllotoxin. CYP3A5 is detected in 10-30% of Caucasian adult livers, and expressed constitutively in adult kidney. CYP3A5 expression does not appear to be sex-influenced and only moderately inducible by xenobiotics both *in vivo* and *in vitro*. CYP3A7 is expressed in fetal liver but only in 25% of adult livers. Molecular mechanisms responsible for the developmentally specific expression of CYP3A's are unknown.

Another supergene family of metabolic enzymes is the glutathione S-transferase (GST) superfamily. These enzymes play an important role in the cellular enzymatic protection against the cytotoxic and mutagenic effects of electrophiles. Thus, GST alleles

associated with impaired detoxification will confer an increased susceptibility to a wide range of diseases. In particular, GST genotypes have been associated with an increased susceptibility in diseases associated with oxidative stress. One of the best characterized of such associations is the null mutation in the mu class GSTM1 gene. GSTM1 is polymorphic due to large deletions in the structural gene. The null GSTM1 genotype is clearly associated with bladder cancer and lung cancer, and possibly associated with colorectal, hepatocellular, gastric, esophageal, head and neck as well as cutaneous cancer. There is considerable evidence that the combination of the GSTM1 null genotype in combination with the cytochrome P450 1A1 rare alleles confers a highly increased risk of developing lung cancers in smokers. The GSTM1 null genotype has also been found to be significantly associated with an increased risk of developing postmenopausal breast cancer as described by Helzlsouer *et al.*, J. Natl. Cancer Inst., 1998, vol. 90, pages 512-518.

Women have a 15 percent lifetime risk of developing breast cancer.

Approximately 10 to 15 percent of all breast cancers are familial, and approximately 33 percent of these may be linked to genetic mutations. Available treatment options include (1) surgery, (2) radiation therapy, (3) chemotherapy, and (4) hormone manipulation. Hundreds of thousands of women are currently undergoing local, as well as systemic, treatment for their breast cancer. Each treatment has its risks and side effects. The risks and side effects of chemotherapy can be substantially reduced and the likelihood of a successful outcome can be increased if the chemotherapeutic regimen selected is tailored to the individual patient. Two drugs commonly used in the treatment of breast cancer are cyclophosphamide and carmustine (BCNU).

Cyclophosphamide is a nitrogen mustard derivative, polyfunctional alkylating agent which is bioconverted from an inert prodrug to an active DNA alkylating agent by the oxidative cytochromes of the liver of which CYP 3A4 and 3A5 are the principle enzymes. Thus, patients with lower levels of either enzyme will produce less of the active forms of cyclophosphamide when given the same dose over the same time as a patient who has normal levels of these activating enzymes. Cyclophosphamide functions

to interfere with DNA replication and transcription of RNA, ultimately resulting in the disruption of nucleic acid function. The drug also exhibits potent immunosuppressive activity and phosphorylating properties that enhance its cytotoxicity. In the treatment of breast cancer, cyclophosphamide used alone has been reported to produce objective
5 responses in about 35% of patients. Used in combination regimens, objective responses have been reported in up to 90% of patients, and cyclophosphamide-containing combinations are believed by some experts to be the treatment of choice.

BCNU is a nitrosourea with a broad spectrum of activity. It is a classic alkylating agent, but also inhibits DNA repair by isocyanate formation. BCNU is used alone or as a
10 component of various chemotherapy regimens in the treatment of primary or metastatic tumors. BCNU is a highly toxic drug with a low therapeutic index, thus a therapeutic response is unlikely without some evidence of toxicity. The primary toxicities are pulmonary toxicity and hepatic dysfunction which appear to be dose related. Patients receiving cumulative doses exceeding 1400mg/m² are at substantially higher risk than
15 patients receiving lower cumulative doses. Thus, any means of screening prospective cancer patients for factors which can effect the dosing of BCNU is of great importance in effectively designing chemotherapy regimens that include BCNU to enhance the clinical outcome while minimizing adverse effects.

Cisplatin is a bifunctional alkylating agent that binds to DNA and inhibits DNA
20 synthesis. The drug produces predominately DNA interstrand crosslinks with some intrastrand crosslinks resulting from the formation of adducts between activated platinum complexes of the drug. Interstrand crosslinking appears to correlate well with the cytotoxicity of the drug. Cisplatin is used to treat a wide variety of neoplasms and is often used as a component of combination chemotherapeutic regimens because of its
25 relative lack of hematologic toxicity. Cisplatin is a highly toxic drug with a low therapeutic index. While hematologic toxicities such as thrombocytopenia and leukopenia are the major dose-limiting adverse effects of cisplatin therapy, other dose-limiting adverse effects including nephrotoxicity, ototoxicity, neurotoxicity, and emesis are frequently seen. These adverse effects are potentiated in patients receiving other

antineoplastic agents or drugs with nephrotoxic or ototoxic effects, such as aminoglycoside antibiotics.

Since the rates of metabolism of drugs and toxins can depend on the amounts and kinds of P450s expressed in a tissue, variation in biological response may be determined
5 by the profile of expression of P450s in each person. As noted above, this variation in response may significantly influence the outcome of treating breast cancer patients with different antineoplastic drugs. Analysis of genetic polymorphisms that lead to altered expression and enzyme activity of these metabolic enzymes are therefore of interest.

SUMMARY OF THE INVENTION

10 The present invention is directed to novel polymorphisms in the human CYP3A4 and CYP3A5 genes and the detection of polymorphisms in genes encoding the CYP3A4, CYP3A5 and GSTM1 metabolic enzymes in individuals. The polymorphisms detected can influence the outcome of breast cancer treatments and the selection of
15 chemotherapeutic agents used to treat breast cancer in the patients tested. These polymorphisms are predictive of altered metabolism of chemotherapeutic agents used in the treatment of breast cancer and/or the occurrence of disease. According to the present invention there are provided CYP3A4 and CYP3A5 polymorphic nucleic acid sequences and methods to use such nucleic acid sequences and polymorphic GSTM1 sequences, in particular for diagnostic purposes to identify individuals having a polymorphic genotype.

20 In one embodiment, the present invention is directed to an isolated nucleic acid molecule, comprising a sequence selected from the sequence of SEQ ID NO:1; and the sequence fully complementary thereto. The present invention is also directed to an isolated nucleic acid molecule which includes at least one base variation from that of the human CYP3A4 sequence, wherein the nucleic acid molecule is selected from a nucleic
25 acid molecule that comprises a G nucleotide for an A nucleotide at position -392 of the promoter of said CYP3A4 gene with respect to the start codon of said CYP3A4 gene and at least 17 other bases of said CYP3A4 gene contiguously appurtenant thereto, and a nucleic acid molecule which is fully complementary thereto. In this embodiment, the nucleic acid molecule can include a sequence selected from the group consisting of SEQ

ID NO:1, and/or a nucleic acid sequence which is fully complementary to SEQ ID NO:1.

In another embodiment, the present invention is directed to an isolated nucleic acid molecule, comprising a sequence selected from the sequence of SEQ ID NO:2; and the sequence fully complementary thereto. The present invention is also directed to an isolated nucleic acid molecule which includes at least one base variation from that of the human CYP3A5 sequence, wherein the nucleic acid molecule is selected from a nucleic acid molecule that comprises a G nucleotide for an A nucleotide at position -147 of the promoter of said CYP3A5 gene with respect to the start codon of said CYP3A4 gene and at least 33 other bases of said CYP3A5 gene contiguously appurtenant thereto, and a nucleic acid molecule which is fully complementary thereto. In this embodiment, the nucleic acid molecule can include a sequence selected from the group consisting of SEQ ID NO:2 and/or a nucleic acid sequence which is fully complementary to SEQ ID NO:2.

A further embodiment of the present invention includes a method of detecting a variant gene having a polymorphism associated with reduced metabolism of a substrate selected from the group consisting of a CYP3A4 substrate, a CYP3A5 substrate and a GSTM1 substrate in an individual. This method includes obtaining a nucleic acid sample comprising a gene isolated from the individual, the gene being selected from a CYP3A4 gene, a CYP3A5 gene and a GSTM1 gene. The method further includes detecting the presence or absence in the individual of a polymorphism selected from (i) a substitution of a G nucleotide for an A nucleotide at position -392 of the promoter of the CYP3A4 gene with respect to the start codon of the CYP3A4 gene, wherein the presence of the substitution is associated with reduced CYP3A4 substrate metabolism: (ii) a substitution of a G nucleotide for an A nucleotide at position -147 of the promoter of the CYP3A5 gene, wherein the presence of the substitution is associated with reduced CYP3A5 substrate metabolism: and (iii) a GSTM1 null mutation, wherein the presence of the GSTM1 null mutation is associated with reduced GSTM1 substrate metabolism. The method can also include determining whether the individual is homozygous or heterozygous for the polymorphism. A variety of assays are suitable for determining whether any of the identified variant sequences are present. The substrates in question for

any one of the three genes can be selected from cyclophosphamide and BCNU.

In a further embodiment, the present invention includes a method for selecting a treatment for a cancer patient. The method the genotyping steps described above and further includes selecting a cancer treatment regime that does not include administration of an anti-cancer agent selected from the group consisting of cyclophosphamide and BCNU if one or more of the polymorphisms are present. A further method for selecting a treatment for a cancer patient can include selecting a cancer treatment regime that includes administration of an anti-cancer agent selected from the group consisting of cyclophosphamide and BCNU if none of the polymorphisms are present.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to nucleic acid molecule compositions that contain certain genetic characteristics and methods that identify the presence or absence of such characteristics in patients. The present invention includes the identification of different genetic polymorphisms in the cytochrome P450 3A4, cytochrome P450 3A5 and the GSTM1 genes. The presence or absence of certain polymorphisms in one or more of these genes has been found to be prognostic for a therapeutic response to certain antineoplastic agents. Identification of these polymorphic sequences is used in individuals to screen for altered metabolism of CYP3A4, CYP3A5, and GSTM1 substrates, potential drug-drug interactions, drug adverse effects, likelihood of successful clinical outcome following treatment with cyclophosphamide, cisplatin and/or BCNU.

It is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, animal species or genera, constructs, and reagents described, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

As used herein the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a construct" includes a plurality of such constructs and reference to "the CYP3A4 nucleic

acid" includes reference to one or more nucleic acids and equivalents thereof known to those skilled in the art, and so forth.

According to the present invention, reference to an "isolated nucleic acid molecule" refers to a nucleic acid molecule which is the size of or smaller than a gene.

5 Thus, an isolated nucleic acid molecule does not encompass isolated genomic DNA or an isolated chromosome. The term isolated nucleic acid molecule does not connote any specific minimum length. It should also be appreciated that reference to an isolated nucleic acid molecule does not necessarily reflect the extent of purity of the nucleic acid molecule. An isolated nucleic acid molecule of the present invention can be obtained
10 from a natural source, such as a tissue sample, or it can be produced using molecular biology techniques, such as by PCR amplification, or it can be produced by chemical synthesis.

"Allele" has the meaning which is commonly known in the art, that is, a genomic variant of a referent gene, including variants, which, when translated result in functional
15 or dysfunctional (including non-existent) gene products. The first identified allelic form is arbitrarily designated as the reference form and other allelic forms are designated as alternative or variant alleles. The allelic form occurring most frequently in a selected population is sometimes referred to as the wildtype form.

"Contiguously appurtenant to" means any bases flanking the referent position,
20 including the instances of all bases selected 5' to the referent position and no bases selected 3' to the referent position; all bases selected 3' to the referent position and no bases selected 5' to the referent position; and some bases selected 5' and some bases selected 3' to the referent position. The term is intended to mean that the selected bases necessarily must be in the same sequential order as described in the referent sequence,
25 with the exception of the variant base at the referent position.

"For the purpose of determining genotype" means that one of the purposes is to determine genotype, not necessarily that the end goal or use of the information is to determine genotype. For instance, "for the purpose of determining genotype" includes the use of the information to determine genotype for the ultimate goal of determining

probability of negative or positive drug interactions.

"Gene" has the meaning that is commonly-known in the art, that is, a nucleic acid sequence that includes the translated sequences that code for a protein ("exons") and the untranslated intervening sequences ("introns"), and any regulatory elements ordinarily
5 necessary to translate the protein.

"Genotype" has the meaning that is commonly-known in the art, that is, a physical description of a nucleic acid sequence.

"Hybridization" has the meaning that is commonly-known in the art, that is, the formation of a duplex structure by two single-stranded nucleic acids due to
10 complementary base pairing. Hybridization can occur between exactly complementary nucleic acid strands or between nucleic acid strands that contain some regions of mismatch.

"Polymorphism" means a polymorphism wherein the group exists by virtue of a difference in identity of one or more nucleotides at given sequence locations. The
15 location of nucleotide identity differences is usually preceded by and followed by highly conserved sequences (e.g., sequences that vary in less than 1/100 or 1/1000 members of the populations). However, more than one single nucleotide polymorphism can exist between or among the group members. A "transition" is the replacement of one purine by another purine or one pyrimidine by another pyrimidine. A "transversion" is the
20 replacement of a purine by a pyrimidine or vice versa. Single nucleotide polymorphisms can also arise from a deletion of a nucleotide or an insertion of a nucleotide relative to a given sequence location.

"Stringent hybridization" means that which is commonly-known in the art, that is, at a salt concentration of no more than 1M and a temperature of at least 25 degrees
25 Celsius. For example, conditions of 5X SSPE (750 mM NaCl, 50 mM Sodium Phosphate, 5 mM EDTA, pH 7.4) and a temperature of 55 degrees to 60 degrees Celsius are suitable.

"Substrate" means a chemical entity that is modified by activity of a CYP3A4, CYP3A5 and/or GSTM1 enzyme, usually under normal physiological conditions. Most of these substrates are lipophilic compounds. Although the duration of drug action tends

to be shortened by metabolic transformation, drug metabolism is not "detoxification." Frequently the metabolic product has greater biologic activity than the drug itself. In some cases the desirable pharmacologic actions are entirely attributable to metabolites, the administered drugs themselves being inert. Likewise, the toxic side effects of some drugs may be due in whole or in part to metabolic products. The range of known substrates for CYP3A4, CYP3A5 and GSTM1 is very broad. In a preferred embodiment, a substrate of any one or more of CYP3A4, CYP3A5 and GSTM1 is selected from the group consisting of cyclophosphamide, cisplatin and BCNU. In a further preferred embodiment, a substrate of any one or more of CYP3A4, CYP3A5 and GSTM1 is selected from the group consisting of cyclophosphamide and BCNU.

All technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs unless clearly indicated otherwise.

In the present invention, alleles are expressed by symbols in accordance with definitions given by IUPAC-IUB and common names or common usage in the art. One embodiment of the present invention is an isolated nucleic acid molecule comprising a CYP3A4 sequence polymorphism of SEQ ID NO:1 as part of other than a naturally occurring chromosome, as described in detail below. Another embodiment of the present invention is an isolated nucleic acid molecule comprising a CYP3A5 sequence polymorphism of SEQ ID NO:2 as part of the other than a naturally occurring chromosome. In accordance with the present invention, an isolated nucleic acid molecule is a nucleic acid molecule that has been removed from its natural milieu (i.e., that has been subject to human manipulation) and can include DNA, RNA, or derivatives of either DNA or RNA. An isolated nucleic acid molecule of the present invention can be isolated from its natural source or can be produced using recombinant DNA technology (e.g., polymerase chain reaction (PCR) amplification, cloning) or chemical synthesis. The nucleic acid molecules of the present invention are isolated and obtained in substantial purity, generally as other than an intact chromosome. Usually, the nucleic acid molecule will be obtained substantially free of other nucleic acid sequences, generally being at

least about 50%, usually at least about 90% pure. Although the phrase "nucleic acid molecule" primarily refers to the physical nucleic acid molecule and the phrase "nucleic acid sequence" primarily refers to the sequence of nucleotides on the nucleic acid molecule, the two phrases can be used interchangeably.

5 The nucleic acid sequence of the CYP3A4 gene is known in the art and accessible in public databases. Genbank Accession No. D11131 provides a partial sequence of the human cytochrome P450 3A4 gene. Genbank Accession No. M18907 (cDNA) provides the cDNA sequence of a human CYP3A4 allele. The promoter region of the CYP3A4 gene is provided as SEQ ID NO:3.

10 The nucleic acid sequence of the CYP3A5 gene is generally known in the art and accessible in public databases. For example, Genbank Accession No. S74699 provides the CYP3A5 5' genomic region. However, the present inventors have sequenced a portion of the promoter region and exon 1 of the CYP3A5 gene and found errors in Genbank Accession No. S74699. The correct sequence for the portion of the promoter and exon 1
15 is provided as SEQ ID NO:4.

 The nucleic acid sequence of the GSTM1 gene is known in the art and accessible in public databases. The null mutation in the mu class GSTM1 gene is well known in the art. For example, see Xu, S.; Wang, Y.; Roe, B.; Pearson, W. R. : Characterization of the human class mu glutathione S-transferase gene cluster and the GSTM1 deletion. J.
20 Biol. Chem. 273: 3517-3527, 1998.

 As used herein, the term "CYP3A4 gene," "CYP3A5 gene" or "GSTM1 gene" is intended to refer to both the wildtype and polymorphic sequences, unless specifically denoted otherwise. Nucleic acids of particular interest comprise the provided polymorphic sequences. It is within the skill of one in the art to identify the location of a
25 polymorphic sequence of the present invention using wildtype CYP3A4, CYP3A5 or GSTM1 genomic or cDNA sequences known in the art. A skilled artisan can use a polymorphic sequence, its corresponding wildtype sequence and the CYP3A4, CYP3A5 or GSTM1 sequence contiguously appertanant to the referenced polymorphism to determine the position of the polymorphism. The positions of the CYP3A4 and CYP3A5

polymorphisms of the present inventions are described below in Table 1.

TABLE 1

GENE	TYPE	SEQ ID NO:	SEQUENCE
CYP3A4	WT	5	GACAAGGGCAAGAGAGAG
	PM	1	GACAAGGGCAGGACAGAG
CYP3A5	WT	6	CGATTCTTTGCTACTGGCTGCAGCTGCAGCCCCA
	PM	2	CGATTCTTTGCTACTGGCTGCAGCTGCAGCCCCG

In accordance with the present invention, the polymorphism of the CYP3A4 sequence occurs at nucleotide -392 of the promotor region of the CYP3A4 genomic sequence. Sequences in the promotor region are denoted by negative numbers corresponding to the number of nucleotides 5' to the start codon. The start codon of the CYP3A4 gene is at nucleotides 1208-1210 of SEQ ID NO:3. The polymorphism is a base pair substitution of A to G. The polymorphism is in the promotor region so it does not result in an amino acid substitution. This CYP3A4 polymorphism is described in detail in WO 9913106A1, published March 18, 1999 and is identified in SEQ ID NO:44 therein. In addition, the CYP3A4 polymorphism can be identified as an A to G transition at position -290 with reference to the transcription initiation site. More particularly, the CYP3A4 polymorphism occurs at nucleotide 816 of SEQ ID NO:3. One embodiment of the present invention is an isolated nucleic acid molecule that comprises the foregoing human cytochrome P450 3A4 polymorphism. Such an isolated nucleic acid molecule includes the polymorphism and at least 17 other bases, alternatively at least 20 other bases, at least 30 other bases, at least 40 other bases, or at least 50 other bases of the wildtype sequence contiguously appurtenant thereto. Further, the present invention includes human CYP3A4 alleles that comprise the CYP3A4 polymorphism as described herein, having appurtenant sequences of 10, 15, 20, 25, 30, 35, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 250, 300, 350, 400, 450, 500, or 1000 bases, or any whole

number encompassed by the range of 1-10,000. Further embodiments of the present invention include nucleic acid molecules having a sequence fully complementary to any of the sequences described above.

5 In accordance with the present invention, the polymorphism of the CYP3A5 sequence occurs at nucleotide -147 of the promotor region of the CYP3A5 genomic sequence. Sequences in the promotor region are denoted by negative numbers corresponding to the number of nucleotides 5' to the start codon. The start codon of the CYP3A5 gene is at nucleotides 1184-1186 of SEQ ID NO:4. The polymorphism is a base pair substitution of A to G. The polymorphism is in the promoter region so it does not result in an amino acid substitution. In addition, the CYP3A5 polymorphism can be identified as an A to G transition at position -44 with reference to the transcription initiation site. More particularly, the CYP3A5 polymorphism occurs at nucleotide 1037 of SEQ ID NO:4. One embodiment of the present invention is an isolated nucleic acid molecule that comprises the foregoing human cytochrome P450 3A5 polymorphism.

10 Such an isolated nucleic acid molecule includes the polymorphism and at least 33 other bases, alternatively at least 40 other bases, or at least 50 other bases of the wildtype sequence contiguously appurtenant thereto. Further, the present invention includes human CYP3A5 alleles that comprise the CYP3A5 polymorphism as described herein, having appurtenant sequences of 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 250, 300, 350, 400, 450, 500, or 1000 bases, or any whole number encompassed by the range of 1-10,000. Further embodiments of the present invention include nucleic acid molecules having a sequence fully complementary to any of the sequences described above.

25 In accordance with the present invention, the polymorphism of the GSTM1 gene is the well-characterized null mutation.

The genotype of an individual is determined with respect to the provided CYP3A4, CYP3A5 and/or GSTM1 gene polymorphisms. The genotype is useful for determining the presence of phenotypically evident polymorphism having the effect of

reduced metabolism of a CYP3A4, CYP3A5 or GSTM1 substrate, respectively.

One embodiment of the present invention is a method of identifying a sample containing a nucleic acid molecule that comprises a wildtype or variant allele, the method comprising identifying the presence or absence of one or more of the CYP3A4, CYP3A5
5 or GSTM1 polymorphisms described herein.

Many of the diagnostic assays rely on amplification of part or all of a CYP3A4, CYP3A5 and/or GSTM1 nucleic acid molecule. In one embodiment, portions of a nucleic acid molecule are amplified by the polymerase chain reaction (PCR). The PCR process is described in e.g., U.S. Pat. Nos. 4,683,195; 4,683,202; and 4,965,188; PCR
10 Technology: Principles and Applications for DNA Amplification (ed. Erlich, Freeman Press, New York, N.Y., 1992); PCR Protocols: A Guide to Methods and Applications (eds. Innis et al., Academic Press, San Diego, Calif. (1990); Mattila et al. Nucleic Acids Res. 19:4967 (1991); Eckert & Kunkel PCR Methods and Applications 1:17 (1991); PCR
15 (eds. McPherson et al., IRL Press, Oxford), each of which is incorporated by this reference in its entirety.

To amplify a portion of a nucleic acid molecule in accordance with the present invention in a sample by PCR, the sequence must be accessible to the components of the amplification system. Accessibility can be achieved by isolating nucleic acid molecules from the sample. A variety of techniques for extracting nucleic acid molecules from
20 biological samples are known in the art. Alternatively, if the sample is fairly readily disruptable, the nucleic acid need not be purified prior to amplification by the PCR technique, i.e., if the sample comprises cells, particularly peripheral blood lymphocytes or monocytes, lysis and dispersion of the intracellular components may be accomplished merely by suspending the cells in hypotonic buffer. See Han *et al.*, Biochemistry, 1987,
25 vol. 26, pages 1617-1625. Polymorphisms are detected in a nucleic acid molecule from an individual being analyzed. For assay of genomic DNA, virtually any biological sample (other than pure red blood cells) is suitable. Examples of convenient tissue samples include whole blood, semen, saliva, tears, urine, fecal material, sweat, buccal, skin and hair. Nucleic acid molecules can be obtained according to procedures well-

known in the art.

For amplification of mRNA sequences, a first step is the synthesis of a DNA copy (cDNA) of the region to be amplified by reverse transcription. Reverse transcription is the polymerization of deoxynucleoside triphosphates to form primer extension products that are complementary to a ribonucleic acid template. The process is effected by reverse transcriptase, an enzyme that initiates synthesis at the 3'-end of the primer and proceeds toward the 5'-end of the template until synthesis terminates. Examples of suitable polymerizing agents that convert the RNA nucleic acid molecule into a complementary, cDNA sequence are avian myeloblastosis virus reverse transcriptase and *Thermus thermophilus* DNA polymerase. Reverse transcription can be carried out as a separate step, or in a homogeneous reverse transcription-polymerase chain reaction (RT-PCR). Polymerizing agents suitable for synthesizing a cDNA sequence from the RNA template are reverse transcriptase (RT), such as avian myeloblastosis virus RT, Moloney murine leukemia virus RT, or *Thermus thermophilus* DNA polymerase.

Primers for PCR amplification are designed so that the position at which each primer hybridizes along a duplex sequence is such that an extension product synthesized from one primer, when separated from the template (complement), serves as a template for the extension of the other primer. The primers are selected to be substantially complementary to the different strands of each specific sequence to be amplified. This means that the primers must be sufficiently complementary to hybridize with their respective strands. Therefore, the primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer with the remainder of the primer sequence being complementary to the strand. Alternatively, complementary bases or longer sequences can be interspersed into the primer, provided that the primer sequence has sufficient complementarity with the sequence of the strand to be amplified to hybridize therewith and thereby form a template for synthesis of the extension product of the other primer. Paired primers for amplification of a given segment of DNA are designated forward and reverse primers. The forward primer hybridizes to a double-stranded DNA molecule at a

position 5', or upstream, from the reverse primer. The forward primer hybridizes to the complement of the coding strand of the double stranded sequence, i.e., the antisense strand, and the reverse primer hybridizes to the coding strand.

5 The appropriate length of a primer depends on the intended use of the primer but typically ranges from about 10 to about 100, preferably about 15 to about 50, more preferably about 15 to about 30, or more preferably about 15 to about 25 nucleotides in length. The spacing of primers determines the length of segment to be amplified. The spacing is not usually critical and amplified segments can range in size from about 25 bases to at least about 35 kilobases in length. Segments from about 25 to about 2000,
10 preferably about 50 to about 1000, more preferably about 100 to about 500 nucleotides in length are typical.

A primer can be labeled, if desired, by incorporating a label detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include ^{32}P , fluorescent dyes, electron-dense reagents, enzymes (as
15 commonly used in an ELISA), biotin, or haptens and proteins for which antisera or monoclonal antibodies are available. A label can also be used to "capture" the primer, so as to facilitate the immobilization of either the primer or a primer extension product, such as amplified DNA, on a solid support.

Other suitable amplification methods include the ligase chain reaction (LCR) (see
20 Wu and Wallace, *Genomics*, 1989, vol. 4, pages 560; Landegren *et al.*, *Science*, 1988, vol. 241, pages 1077; transcription amplification (Kwoh *et al.*, *Proc. Natl. Acad. Sci. USA*, 1989, vol. 86, page 1173), and self-sustained sequence replication (Guatelli *et al.*, *Proc. Nat. Acad. Sci. USA*, 1990, vol. 87, page 1874) and nucleic acid based sequence amplification (NASBA). The latter two amplification methods involve isothermal
25 reactions based on isothermal transcription, which produce both single stranded RNA (ssRNA) and double stranded DNA (dsDNA) as the amplification products in a ratio of about 30 or 100 to 1, respectively.

An allele-specific primer can be used in a PCR amplification. The allele-specific primer hybridizes to a site on a nucleic acid molecule that overlaps with a polymorphism

and extension will only occur if an allelic form complementary to the primer is present. See Gibbs, Nucleic Acid Res., 1989, vol. 17, pages 2427-2448. This primer is used in conjunction with a second primer which hybridizes at a distal site. Amplification proceeds from the two primers leading to a detectable product signifying the particular allelic form is present. Thus, the presence or absence of an amplification product is detected using standard methods. Controls can be used that test the efficacy of the amplification reaction itself or that allow the experimental results to be compared with known wildtype or polymorphic CYP3A4, CYP3A5 and/or GSTM1 nucleic acid molecule samples. The method works best when the mismatch is included in the 3'-most position of the oligonucleotide aligned with the polymorphism because this position is most destabilizing to elongation from the primer.

Sample nucleic acid molecules, isolated directly from cells, amplified or cloned fragments, can also be analyzed by a number of other methods known in the art. The nucleic acid molecule can be sequenced by using either the dideoxy chain termination method or other methods (see for example Sambrook *et al.*, Molecular Cloning, A Laboratory Manual (2nd Ed., CSHP, New York 1989); Zyskind *et al.*, Recombinant DNA Laboratory Manual, (Acad. Press, 1988)).

Hybridization using allele-specific probes, described by e.g., Saiki *et al.*, Nature 324, 163-166 (1986); Dattagupta, EP 235,726, Saiki, WO 89/11548, can be used to determine the presence or absence of a polymorphism by, for example Southern blot, dot blots, etc. An allele-specific probe can be designed that hybridizes to a segment of a nucleic acid molecule from one individual but does not hybridize to the corresponding segment from another individual due to the presence of different polymorphic forms in the two individuals. Hybridization conditions should be sufficiently stringent that there is a significant difference in hybridization intensity between alleles.

The hybridization pattern of a control and variant sequence to an array of oligonucleotide probes immobilized on a solid support, as described in US Pat. No. 5,445,934, or in WO 95/35505, can also be used as a means of detecting the presence of variant sequences.

Amplification products generated using the polymerase chain reaction can be analyzed by the use of denaturing gradient gel electrophoresis (DGGE). Different alleles can be identified based on the different sequence-dependent melting properties and electrophoretic migration of DNA in solution. Erlich, ed., PCR Technology, Principles and Applications for DNA Amplification, (W.H. Freeman and Co, New York, 1992), Chapter 7.

Alleles of target sequences can be differentiated using single-strand conformation polymorphism analysis (SSCP), which identifies base differences by alteration in electrophoretic migration of single stranded PCR products, as described in Orita *et al.*, Proc. Nat. Acad. Sci. 86, 2766-2770 (1989). Amplified PCR products can be generated as described above, and heated or otherwise denatured, to form single stranded amplification products. Single-stranded nucleic acids may refold or form secondary structures which are partially dependent on the base sequence. The different electrophoretic mobilities of single-stranded amplification products can be related to base-sequence difference between alleles of target sequences.

Other methods of detection include mismatch cleavage detection and heteroduplex analysis in gel matrices. These methods are used to detect conformational changes created by DNA sequence variation as alterations in electrophoretic mobility. Alternatively, where a polymorphism creates or destroys a recognition site for a restriction endonuclease, referred to as restriction length polymorphism, or RFLP, the sample is digested with that endonuclease and the products size fractionated to determine whether the fragment was digested. Fractionation is performed by gel or capillary electrophoresis, particularly acrylamide or agarose gels.

In one embodiment of the present invention, an array of oligonucleotides are provided, where discrete positions on the array are complementary to one or more of the provided polymorphic sequences, e.g. oligonucleotides of at least 12 nucleotides, frequently 20 nucleotides or larger and including the sequence flanking the polymorphic position. Such an array may comprise a series of oligonucleotides, each of which can specifically hybridize to a different polymorphism. For examples of arrays, see Hacia *et*

al., 1996, Nat. Genet., vol. 14, pages 441-447 and DeRisi *et al.*, 1996, Nat. Genet., vol. 14, pages 457-460. Arrays of interest may further comprise sequences, including polymorphisms, of other genetic sequences, particularly other sequences of interest for pharmacogenetic screening.

5 It is within the scope of the present invention that one or more CYP3A4, CYP3A5 and GSTM1 polymorphisms provided herein can be detected in a single assay such as a multiplex assay to identify the presence or absence of different alleles in the same assay, see for example Stuvén *et al.*, Pharmacogenetics, 1996, vol. 6, pages 417-421.

10 According to the present invention, a polymorphism provided herein is indicative of a poor clinical outcome of breast cancer treatment using cyclophosphamide, BCNU or combinations of these drugs together or in combination with other chemotherapeutic regimens.

15 A preferred strategy for analysis entails amplification of a DNA sequence spanning the polymorphism, in the instance of CYP3A4 or CYP3A5 polymorphisms of the present invention. Amplification of such a sequence can be primed from forward and reverse primers that hybridize to a CYP3A4 or CYP3A5 gene on opposite sides of the polymorphism but which do not hybridize to the variant nucleotide sequence itself. That is, for detection of the polymorphism, the forward primer hybridizes upstream or 5' to the polymorphic site and the reverse primer hybridizes downstream or 3' to this site. The
20 forward primer is sufficiently complementary to the antisense strand of a nucleic acid molecule to hybridize therewith and the reverse primer is sufficiently complementary to the sense strand of the sequence to hybridize therewith. The primers usually comprise first and second subsequences from opposite strands of a double-stranded DNA sequence. It is particularly important to avoid mismatches in the two nucleotides at the 3' end of the
25 primer (especially the terminal nucleotide). Having amplified a segment of a gene known to span a polymorphism, a variety of assays are available for determining whether the polymorphism is present that are disclosed herein, preferably, using allele specific primers. For example, selective amplification of the wildtype allele of the gene in question allele can be accomplished using a forward primer that has about 10-50, and

usually 15-30 nucleotides from the wildtype genomic sequence, including the portion of the gene corresponding to the position of the polymorphism. Such a forward primer when paired with any suitable reverse primer downstream from the polymorphic nucleotide sequence (i.e., sufficiently complementary to the sense strand of gene to hybridize therewith) can be used to amplify selectively the wildtype allele without amplifying a mutant allele. The polymorphic nucleotide sequence usually occurs near, or preferably at, the 3' end of the primer. The same result can be achieved by using a reverse primer that has about 10-50 or usually 15-30 contiguous nucleotides from the complement of the wildtype genomic sequence (i.e., the antisense strand) including the polymorphic nucleotide sequence. Such a reverse primer can be paired with any suitable forward primer sufficiently complementary to a sequence of the antisense strand of the gene upstream from the polymorphic nucleotide sequence to hybridize therewith. The position corresponding to the polymorphic nucleotide sequence should again be at or near the 3' end of the reverse primer. For selective amplification of a mutant allele a suitable forward primer for amplification comprises about 10-50 or usually 15-30 contiguous nucleotides including a polymorphic nucleotide sequence from the mutant genomic sequence (i.e., the sense strand). The forward primer can be paired with any suitable reverse primer sufficiently complementary to the sense strand of a genomic subsequence downstream from the polymorphic nucleotide sequence to hybridize therewith. Alternatively, the same result can be achieved using a reverse primer comprising about 10-50 or 15-30 contiguous nucleotides including the polymorphic nucleotide sequence from the complement of the mutant sequence (i.e., the antisense strand). Such a reverse primer can be paired with any suitable forward primer sufficiently complementary to the antisense strand of a subsequence upstream from a polymorphic nucleotide sequence to hybridize therewith.

Following amplification, the sample under test is characterized as wildtype or mutant by the presence or absence of an amplification product. With a primer designed for selective amplification of the wildtype allele, the presence of an amplification product is indicative of that allele and the absence of an amplification product indicative of a

mutant allele. The converse applies for primers designed for selective amplification of a mutant allele. In preferred assay, a sample is divided into two aliquots, one of which is amplified using primers for wildtype allele amplification, the other of which is amplified using primers appropriate for mutant allele amplification. The presence of an
5 amplification product in one but not both of the aliquots indicates that the individual under test is either wildtype or a homozygous for the mutation (depending on aliquot in which the amplification product occurred). The presence of amplification product in both aliquots indicates that the individual is heterozygous. The absence of an amplification product in both aliquots would indicate either the absence of a gene or a quality control
10 problem in the amplification procedure requiring that the assay be repeated. The presence or absence of amplification products can be detected by gel electrophoresis using methods standard in the art or described herein.

In the case of the GSTM1 null mutation, a preferred strategy for analysis entails amplification using any portion of sequence in the coding region of the GSTM1 gene that
15 would produce a detectable PCR product if the gene is wildtype, but no PCR product if the null mutation is present. Although some conventional methods for determining hetero- or homozygosity, such as that described above, would not be useful for the GSTM1 null mutation, those skilled in the art could design suitable methods. For example, one could conduct an assay in which two differently sized PCR amplification
20 products would be produced for a heterozygote. Such an assay could be designed once the breakpoints for the null mutation were determined.

In further preferred embodiments of the present invention, the various genotyping methods described herein can be used in a method for selecting a treatment for a cancer patient. As described below in the examples, the polymorphisms of the present invention
25 have been associated with reduced metabolism of various chemotherapeutic agents, specifically cyclophosphamide and BCNU. Thus, in such a method for selecting a treatment for a cancer patient, if one or more of the polymorphisms are present, a cancer treatment regime is selected to account for the phenotypic result of the identified genotype. More particularly, a cancer treatment regime can be selected that does not

include administration of an anti-cancer agent that is metabolized by an expression product of one of the genes described herein, and particularly, cyclophosphamide and BCNU. Alternatively, a cancer treatment regime can be selected that includes administering an anti-cancer agent that is metabolized by an expression product of one of the genes described herein at higher than conventional doses. More particularly, such doses can be at least about 1% higher than conventional doses, more preferably, at least about 10 % higher than conventional doses, and even more preferably, at least about 50 % higher than conventional doses. Conventional doses of known chemotherapeutic agents are well-known to those of skill in the art. More particularly, conventional doses of cyclophosphamide can include 40-100 mg/Kg. Conventional doses of BCNU can include 100-200 mg/m² every 6 weeks. In an alternative method for selecting a treatment for a cancer patient using one of the various genotyping methods described herein, if the patient is identified as having none of the three polymorphisms of the present invention, a cancer treatment regime that includes administration of an anti-cancer agent selected from the group consisting of cyclophosphamide and BCNU is selected. More particularly, this method includes administration of one of such anti-cancer agents at conventional doses.

One embodiment of the present invention is a diagnostic kit. The kit comprises useful components for practicing the methods of the present invention. The kit typically comprises at least one of the primers needed for the PCR amplification if PCR amplification is used and also control DNA suitable for determining the success of the PCR reaction and/or to confirm the identification of the presence or absence of a polymorphism in a sample. A kit usually contains a matched pair of forward and reverse primers as described above for amplifying a segment encompassing a polymorphism of the present invention. For selective amplification of mutant or wildtype alleles, kits usually contain a pair of primers for amplification of the mutant allele and/or a separate pair of primers for amplification of the wildtype allele. Optional additional components of the kit include, for example, restriction enzymes for analysis of amplification products, reverse-transcriptase or polymerase, the substrate nucleoside triphosphates, and the

appropriate buffers for reverse transcription, PCR, or restriction enzyme reactions. Usually, the kit also contains instructions for carrying out the methods.

5 The method of the present invention is characterized by detecting the polymorphisms provided herein, and is useful in gene diagnosis for detecting CYP3A4, CYP3A5 and/or GSTM1 gene polymorphisms. As long as the method is capable of detecting the aforementioned specific types of mutation which are clearly defined and characterized by the present invention, no limitation is imposed on the techniques to be employed in the method. For example, a variety of routine methods may be widely used. Since the types of gene mutation to be detected by the present invention are now clarified and specified, it would be obvious for skilled persons in the art to adopt another suitable method. For detecting them from the reading of the disclosure of this specification.

Also provided are databases comprising sequence information pertaining to at least one the polymorphisms of the present invention. Software programs comprising use of a database of the present invention are also included.

15 In another embodiment, said method further comprises digesting DNA encoding at least a part of the nucleic acid sequence containing the polymorphism with a restriction enzyme that will cut, or will not cut, at or adjacent to one of the polymorphic positions according to whether the polymorphism is present. Those methods wherein said nucleic acid is amplified prior to the digestion step are preferred. The materials useful for these methods can be obtained as described, and these methods can be accomplished as discussed.

20 Polyclonal and/or monoclonal antibodies that specifically bind to variant gene products but not to corresponding prototypical gene products are also provided. Antibodies can be made by injecting mice or other animals with the variant gene product or synthetic peptide fragments thereof. Monoclonal antibodies are screened as are described, for example, in Harlow & Lane, Antibodies, A Laboratory Manual, Cold Spring Harbor Press, New York (1988); Goding, Monoclonal antibodies, Principles and Practice (2d ed.) Academic Press, New York (1986). Monoclonal antibodies are tested for specific immunoreactivity with a variant gene product and lack of immunoreactivity to

the corresponding prototypical gene product. These antibodies are useful in diagnostic assays for detection of the variant form, or as an active ingredient in a pharmaceutical composition.

5 The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention.

EXPERIMENTAL

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the subject invention, and are not intended to limit the scope of what is regarded as the invention.

10 Efforts have been made to ensure accuracy with respect to the numbers used (e.g. amounts, temperature, concentrations, etc.) but some allowance should be made for experimental errors and deviations. Unless otherwise indicated, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees centigrade; and pressure is at or near atmospheric.

15 Example 1

This example describes the identification of a variant of the known cytochrome P450 3A5 sequence (CYP4503A5).

Blood specimens from 32 individuals were collected after obtaining informed consent. All samples were stripped of personal identifiers to maintain confidentiality.

20 The only data associated with the sample were self-reported gender and racial group designations. Of the 32 individuals, 10 were African Americans, 10 were Caucasians, 6 were Japanese and 6 were Chinese. Genomic DNA was isolated using standard methods. Polymerase chain reaction (PCR) amplification of regions of the CYP4503A5 gene were performed using the primers listed in Table 2. The PCR amplification was performed in

25 a total reaction volume of 50 microliters (μl). The final magnesium chloride concentration (2mM) was optimized empirically. The final genomic DNA concentration was about 100 nanogram (ng) per reaction from 2 individuals. The PCR reactions were performed using Perkin Elmer GeneAmp PCR kit (available from Perkin Elmer, Norwalk, CN) using Taq Gold DNA polymerase according to manufacturer's instructions

and using the following primers.

Table 2. PCR Primers

Region	Forward/Reverse	SEQ ID NO:	5'-3'
3A5	Forward	7	GCAGGTCATTATGTTAGGT
3A5	Reverse	8	CCTTCTTCAACTGTCTCCT

Thermal cycling was performed with an initial denaturation step at 94°C for 10 min, followed by 40 cycles of denaturation at 95°C for 30 sec, primer annealing at 55°C for 45 sec, and primer extension at 72°C for 2 min, followed by final extension at 72°C for 5 min.

The resulting PCR products were purified using Microcon-100 columns (available from Millipore, Bedford, MA). PCR products from two individuals were combined for each cycle of sequencing. Cycle sequencing was performed on the GeneAmp PCR System 9600 PCR machine using the ABI Prism dRhodamine Terminator Cycle Sequencing Ready Reaction Kit (available from Applied Biosystems, Inc., Foster City, CA) according to the manufacturer's directions. Oligonucleotide primers used for the sequencing reactions include those shown in Table 3.

Table 3. Sequencing Primers

Region	Forward/Reverse	SEQ ID NO:	5'-3'
3A5	Forward	9	GCAGGTCATTATGTTAGGT
	Forward	10	CTGATACATAGTTATCTTCCTTG
	Forward	11	CAAGTCAACTCCACCAAC
	Forward	12	GGGCACAAGTACACTCC
	Reverse	13	AACATAGATGAAGAGACTTACC TG

	Reverse	14	CTAAGGGCACAGTCTGG
	Reverse	15	TTCCAGAATACTTGAAATCC
	Reverse	16	TGTGCTGTTGTTTGCTG

5 About 8 µl sequencing reactions were subjected to 30 cycles at 96°C for 20 sec, 50°C for 20 sec, and 60°C for 4 min, followed by ethanol precipitation. Samples were evaporated to dryness at 50°C for about 15 min and resuspended in 2 µl of loading buffer (5:1 deionized formamide:50 mM EDTA pH 8.0), heated to 65°C for 5 min, and electrophoresed through 4% polyacrylamide/6M urea gels in an ABI 377 Nucleic Acid Analyzer according to the manufacturer's instructions for to obtain sequence information. All sequences were determined from both the 5' and 3' (sense and antisense) direction. The 16 electropherograms were analyzed by comparing peak heights, looking for about 25% reduction in peak size and/or presence of extra peaks as an indication of heterozygosity.

15 A sequence comprising a polymorphism identified from the sequencing is shown below the corresponding wildtype sequence, both the polymorphic nucleotide and the corresponding wildtype nucleotide at the same position are shown in bold, and are listed below in Table 4. For example, a variation of a A to a G transition was discovered at base pair -147 in the promoter region of the CYP4503A5 gene.

20 **Table 4.** Newly Identified CYP4503A5 Gene Polymorphism

Location	SEQ ID NO	Polymorphism Sequence
Promoter -147	6	CGATTCTTTGCTACTGGCTGCAGCTGCAGCCCCA
	2	CGATTCTTTGCTACTGGCTGCAGCTGCAGCCCCG

25

SEQ ID NO:4 lists the sequence of the reference CYP4503A5 gene, includes a portion of

the promoter and exon 1.

Example 2

This example describes the association of the Promoter -392 CYP4503A4 polymorphism with survival rate of breast cancer patients receiving treatment with the chemotherapy drugs cyclophosphamide, cisplatin and 1,3-bis-(chloroethyl)-1-nitrosourea.

Genomic DNA was isolated from peripheral blood lymphocytes of 86 chemotherapy naive female patients with metastatic or inflammatory breast cancer who participated in a trial for high-dose cyclophosphamide (CY), cisplatin and 1,3-bis-(chloroethyl)-1-nitrosourea (BCNU) chemotherapy with breast cancer using standard methods. DNA from each patient was genotyped for the Promoter -392 CYP4503A4 single nucleotide polymorphisms (SNPs) in the CYP3A4 gene. Taqman assays were performed using DNA samples from each individual to identify the presence or absence of the Promoter -392 CYP4503A4 variant. The primers identified in Table 5 were used:

Table 5. PCR Primers

PCR Primer	SEQ ID NO:	Primer Sequence
3A4-392 forward	17	ATCTGTAGGTGTGGCTTGTTGG
3A4-392 reverse	18	TATCAGAACTCAAGTGGAGCCAT

PCR amplification was performed using 1X Perkin-Elmer Taqman Reagent Mix #43C4447, about 900 nM 3A4-392 for primer, about 900 nM 3A4-392 rev primer, about 200 nM normal FAM-labeled probe having the nucleic acid sequence 5'-AGAGACAAGGGCAAGAGAGAGGCGAT-3' (SEQ ID NO:19), and 200 nM variant

TET-labeled probe having the nucleic acid sequence 5'-

GACAAGGGCAGGAGAGAGGCGA-3' (SEQ ID NO:20). Thermal cycling was

performed at an initial temperature of 50°C for 2 min followed by a denaturation step at

95°C for 10 min, followed by 40 cycles of denaturation at 94°C for 30 sec, primer

5 annealing and extension was performed at 60°C 30 sec. The fluorescence resulting from the release of labeled probe during PCR amplification and probe hybridization was measured using a fluorometer and the ratio of FAM to TET fluorescence was calculated to determine the occurrence of the polymorphic site and homo- or heterozygosity compared to sequenced controls.

10 The results correlating survival with treatment are shown below in Table 6.

Patients carrying the variant for CYP3A4 demonstrated significantly higher blood

concentrations of CY on day 2 (p=0.0260) and day 3 (p=0.0247). Apparent induction of

CY metabolism (Day 3 exposure/Day 1 exposure) was also higher among those carrying the variant SNP (p=0.0260). Median survival for patients carrying the promoter -392

15 CYP3A4 polymorphism was about 1.6 years compared to about 2.9 years for patients without the variant.

Table 6. CYP4503A4 Variants and Survival. The A allele is wildtype and the B allele is mutant.

Survival >3 years	AA	AB or BB	Total
No	31 (48.3%)	11 (84.6%)	42
Yes	27(51.3%)	2(15.44)	29
Total	58	13	71

5

Mutation Carrier	Median Survival Rate (Years)	95% Confidence Interval (years)	
		Lower Bound	Upper Bound
No	2.88	1.89	4.42
Yes	1.64	0.88	2.43
p-value	0.0476		

10

The results indicate that persons who are homozygous for the A allele have lower blood concentrations of CY and subsequently a 46.6% survival rate beyond 3 years compared to a 15.4% survival rate for those women who have the B allele. Thus, breast cancer patients who have the B allele have higher concentrations of CY and about a 1/3 chance of survival when treated with CY than patients that are homozygous for the A allele. The data suggests that the Promoter -392 variation is detrimental to successful treatment with CY and patients with tumors for which CY treatment is typically used can be genotyped, i.e. screened, as candidates for treatment with CY. Those patients with the B allele, either AB or BB genotypes, will be less likely to benefit from treatment with CY.

15

20

Example 3

This example describes the association of the Promoter -147 CYP4503A5 polymorphism described in Example 2 with survival rate of breast cancer patients

receiving treatment with the chemotherapy drugs CY, cisplatin and BCNU.

The genomic DNA described above in Example 2 was genotyped for the Promoter -147 CYP4503A5 single nucleotide polymorphisms (SNPs) in the CYP3A5 gene.

Taqman assays were performed using DNA samples from each individual to identify the presence or absence of the Promoter -147 CYP4503A5 variant The primers described in Table 7 were used:

Table 7. PCR Primers

PCR Primer	SEQ ID NO:	Primer Sequence
3A5-147 forward	21	GGTGTGTGCGATTCTTTGC
3A5-147 reverse	22	CCCTGCACAGCAGTCTTAGG

PCR amplification was performed using 1X Perkin-Elmer Taqman Reagent Mix

#43C444 7 with about 900 nM 3A5-147 for primer, about 900 nM 3A5-147 rev primer, about 150 nM FAM-labeled probe having the nucleic acid sequence 5'-

CTGCAGCCCCACCTCCTTCTCC-3' (SEQ ID NO:23) and 250 nM variant VIC-

labeled probe having the nucleic acid sequence 5'-CTGCAGCCCCGCCTCCTTCTC-3'

(SEQ ID NO:24). Thermal cycling was performed with at initial temperature of 50°C for

2 min for activation of the Amperase UNG in the Taqman Reagent Mix followed by a

denaturation step at 95°C for 10 min, followed by 40 cycles of denaturation at 60°C for 30

sec, primer annealing and extension was performed at 60°C for 30sec. The fluorescence

resulting from the release of probe labels during PCR and probe hybridization was

measured using a fluorometer and the ratio of FAM to VIC fluorescence was calculated

to determine the occurrence of the polymorphic site and homo- or heterozygosity compared to sequenced controls.

The results correlating survival with treatment are shown below in Table 8.

Patients carrying the variant for CYP3A5 demonstrated significantly higher blood

5 concentrations of CY on day 2 blood concentrations of CY were increased ($p=0.0281$).

These patients also showed increased blood concentrations of BCNU ($p=0.0479$).

Median survival for patients carrying the CYP3A5 SNP was about 1.3 years compared to about 2.9 years for patients without the variant.

Table 8. CYP4503A5 Variants and Survival. The A allele is wildtype and the B allele is
10 mutant.

Survival >3 years	AA	AB or BB	Total
No	35(51.8%)	12(85.7%)	47
Yes	32(48.2%)	2(14.3%)	34
Total	67	14	81

15

Mutation Carrier	Median Survival Rate (Years)	95% Confidence Interval (Years)	
		Upper Bound	Lower Bound
No	2.91	2.28	4.42
Yes	1.33	0.75	2.76
20 p-value	0.0261		

The results indicate that persons who are homozygous for the A allele have a 48.2%

survival beyond 3 years compared to a 14.3% survival rate for those women who have the B allele. Blood concentrations of CY were significantly higher in patients with the B alleles and subsequently these patients showed less than 1/3 chance of survival when treated with CY than patients that are homozygous for the A allele. The data suggests that the Promoter -147 variation is detrimental to successful treatment with CY and patients with tumors for which CY treatment is typically used can be genotyped, i.e. screened, as candidates for treatment with CY. Those patients with the B allele, either AB or BB genotypes, will be less likely to benefit from treatment with CY and should be either given more of the drug, or an alternative chemotherapeutic agent with or without CY.

Example 4

This example describes the association of the null GSTM1 polymorphism with survival rate of breast cancer patients receiving treatment with the chemotherapy drugs cyclophosphamide, cisplatin and BCNU

The genomic DNA described above in Example 2 was genotyped for the null mutation of the GSTM1 gene described in detail in Abdel-Rahman *et al.*, Cancer Letters, vol. 107, pages 229-233, 1996. Fluorescent PCR assays were performed using DNA samples from each individual to identify the presence or absence of the GSTM1 null variant. The primers described in Table 9 were used:

Table 9. PCR Primers

PCR Primer	SEQ ID NO:	Primer Sequence
GSTM1-null forward	23	GAACTCCCTGAAAAGCTAAAGC
GSTM1-control forward	24	GAACTGCCACTTCAGCTGTCT
GSTM1-null reverse	25	GTTGGGCTCAAATATACGGTGG
GSTM1-control reverse	26	CAGCTGCATTTGGAAGTGCTC

PCR amplification was performed at a $MgCl_2$ concentration of 2.5 mM in 1 X buffer D (33.5 mM Tris-HCl, pH 8, 8.3 mM $(NH_4)_2SO_4$, 25 mM KCl and 0.085 mg/ml BSA) with 1.25 mM dNTPs, and 0.05 U/ μ l Perkin Elmer AmpliTaq Gold polymerase. The following primers were used in the PCR reaction: about 300 nM TET-labeled GSTM1-null for primer, about 300 nM unlabeled GSTM1-null rev primer, about 300 nM HEX-labeled GSTM1-control for primer and about 300 nM unlabeled GSTM1-control rev primer. Thermal cycling was performed with an initial denaturation step at 94°C for 5 min, followed by 35 cycles of denaturation at 94°C for 2 min, primer annealing at 59°C for 1 min and extension at 72°C for 1 min, followed by 10 min at 72°C. The resulting PCR products were resolved using standard acrylamide gel electrophoresis methods. The fluorescence emitted from the GSTM1 PCR products and the control PCR products were compared using ABI Prism GeneScan 2.1 software (available from ABI).

The results correlating survival with treatment are shown below in Table 10. Patients homozygous for the GSTM1 null mutation had lower blood concentrations of BCNU ($p=0.0240$). These patients also showed improved tumor response with 58% showing complete response to therapy while only 42% of patients with both copies of the

gene showed complete response ($p=0.043$).Median survival for patients with both GSTM1 deletions was 3.8 years compared to 1.8 years for patients with one or both GSTM1 copies.

Table 10. GSTM1 null mutation and Survival. Positive (+) indicates at least one copy of GSTM1, negative (-) indicates both copies are deleted.

5

un-
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210

Survival >3 years	0	-	Total
No	32(70.3%)	16(44.4%)	48
Yes	13(29.7%)	20(55.6%)	33
Total	45	36	81

5

At Least one GSTM1 gene	Median Survival Rate (Years)	95% Confidence Interval (years)	
		Lower Bound	Upper Bound
No	3.84	2.43	--
Yes	1.81	1.43	2.87
p-value	0.0109		

10

The results indicate that persons who are missing both copies of GSTM1 have a 55.6% rate of survival beyond 3 years compared to a 29.7% survival rate for those women who have one or both copies of GSTM1. The patients missing both copies of GSTM1 had lower blood concentrations of BCNU and subsequently had a 2-fold increased chance of survival when treated with BCNU than patients with one of both copies of GSTM1. Patients with tumors for which BCNU treatment is typically used, can be genotyped, i.e. screened, as candidates for treatment with BCNU. These data suggest that absence of the GSTM1 genes may be beneficial in effective treatment with BCNU.

15

20

While various embodiments of the present invention have been described in detail, it is apparent that modifications and adaptations of those embodiments will occur to those skilled in the art. It is to be expressly understood, however, that such modifications and adaptations are within the scope of the present invention, as set forth in the following claims.